

Two-Electron Three-Centered Bond in Side-On (η^2) Uranyl(V) Superoxo ComplexesVyacheslav S. Bryantsev,^{*,†} Wibe A. de Jong,[‡] Kevin C. Cossel,[†] Mamadou S. Diallo,[†] William A. Goddard III,^{*,†} Gary S. Groenewold,[§] Winnie Chien,^{||} and Michael J. Van Stipdonk^{||}

Materials and Process Simulation Centre, Beckman Institute 139-74, California Institute of Technology, Pasadena, California 91125, William R. Wiley Environmental Molecular Sciences Laboratory, Pacific Northwest National Laboratory, P.O. Box 999, Richland, Washington 99352, Department of Chemical Sciences, Idaho National Laboratory, 2525 North Fremont Avenue, Idaho Falls, Idaho 83415, and Department of Chemistry, Wichita State University, 1845 Fairmount, Wichita, Kansas 67260

Received: May 12, 2008

Theoretical calculations suggest a novel two-electron three-atom bonding scheme for complexation of O₂ with U(V) compounds, leading to the stabilization of superoxo complexes in the side-on (η^2) configuration. This binding motif is likely to play an important role in the oxidative processes involving *trans*-uranium compounds having valence 5f_q electrons.

Introduction

Mononuclear dioxygen–metal compounds, such as FeO₂ complexes with Schiff base and porphyrin ligands, play an essential role in the chemistry of an array of molecules including oxyhemoglobin, cytochrome P-450, and cytochrome oxidase. It is well-known that the superoxo complexes of these molecules have end-on (η^1) coordination geometries with O–O bond lengths of ~ 1.30 Å, Fe–O–O bond angles of $\sim 135^\circ$, and vibrational frequencies of ~ 1150 cm⁻¹,¹ which reflect a formal change of the oxidation state from Mⁿ⁺ to Mⁿ⁺¹. In addition, there are side-on (η^2) peroxo-species, such as [Fe(porphyrin)(O₂)]⁻, in which the O–O bond lengths are ~ 1.46 Å with vibrational frequencies of ~ 820 cm⁻¹.¹ These reflect a formal change of the oxidation state from Mⁿ⁺ to Mⁿ⁺².

We report here computed structures for dioxygen metal compounds in which O₂ is strongly bound side-on (η^2), as it would be in a peroxide but with an O–O bond length of ~ 1.30 Å and frequency of ~ 1200 cm⁻¹, as it would in a superoxide. The systems exhibiting this behavior are dioxouranium(V) complexes in which strongly bound 5f electrons are available for binding of O₂. Addition of O₂ is consistent with condensed-phase studies that suggested that dioxygen complexes are intermediate in the oxidation of U(V) species.² Recently, gas-phase studies have been reported in which O₂ adds to [UO₂(L)_n]⁺ coordination complexes, where L are σ -donor ligands.³ Serial dissociation reactions of the resulting [UO₂(L)₃(O₂)]⁺ complex showed an unusual pattern with initial elimination of neutral L, followed by loss of O₂, even though there were still σ -donors available for elimination. It was hypothesized that dioxygen was oxidizing the U(V) center, forming a U(VI)–superoxide complex whose formation and

stability was in some way influenced by the number of donor ligands. However, these earlier studies had no direct experimental structural data supporting this idea. Here, we employ quantum chemical calculations to understand these spectroscopic observations.

To probe the nature and magnitude of these interactions, we used the relativistic effective core potential (RECP) density functional theory (DFT)⁴ to calculate the geometries, frequencies, and electronic binding energies for 1:1 complexes of U(VI), U(V), and U(IV) dioxo monocations with dioxygen in the two binding motifs (Figure 1 and Table 1). This does not include spin–orbit effects, which we expect not to alter the conclusions of this study.⁵

We first consider the UO₂⁺ cation, which has strong σ and π bonds to both axial oxygen atoms leaving a single electron in one of the 5f_q orbitals (² Φ state).⁶ Dioxygen binding in an η^2 fashion (complex **1**) results in a strong bond (19.4 kcal/mol) between the singly occupied uranyl 5f_q orbital and the O₂ π^* orbital in the equatorial plane (π^*_{xy}) leaving the second π^* singly occupied (π^*_{yz}). The O–O bond length of 1.29 Å and O–O stretching frequency of 1243 cm⁻¹ in complex **1** are consistent with formal reduction of coordinated dioxygen to superoxide.¹

It has been suggested³ that side-on binding of O₂ to U(V) complexes might occur through coupling uranyl 5f_q with the dioxygen π^*_{yz} orbital (perpendicular to the equatorial plane). However, we find that this coupling leads to a bond strength of ~ 4 kcal/mol and requires a 5f_q→5f_q promotion energy of ~ 5 kcal/mol. Thus, these orbitals do not participate in bonding. We were also able to find a stable end-on complex **2**, but this was bound by only 8.5 kcal/mol at the B3LYP level.

To investigate whether these results depend on the flavor of the DFT theory, we calculated the relative bond energies for the side-on coordination geometry, **1**, relative to the end-on, **2**. In all cases, complex **1** appears to be the most stable, by 17.6 (LDA), 15.4 (PBE), 15.5 (PW91), and 10.9 kcal/mol (B3LYP).

* Corresponding author. Phone: 626 395 2730. Fax: 626 585 0918. E-mail: wag@wag.caltech.edu.

[†] California Institute of Technology.

[‡] Pacific Northwest National Laboratory.

[§] Idaho National Laboratory.

^{||} Wichita State University.




Figure 1. Structures and binding energies (kcal/mol) for ground-state dioxygen–uranium complexes 1–8. The weaker interaction is indicated by the thinner dashed line.

TABLE 1: Electronic Binding Energies (BE), Selected Geometric Parameters, and Frequencies for Complexes 1–8^a

system	sym	state	BE kcal/mol	U··O ₂ Å ^c	O–O Å	$\nu_{\text{O–O}}$ cm ⁻¹
1	C_{2v}	2A_2	19.4	2.293	1.290	1243
		${}^2A_1'$	8.5	2.613	1.205	1609
2	C_s	${}^4A_1'$	7.7	2.717	1.203	1637
		2A_1	25.6	2.318	1.298	1233
3	C_1	2A	29.3	2.338	1.305	1225
		2A	28.0	2.369	1.301	1240
4	C_s	${}^1A_1'$	36.3 ^b	2.238	1.318	1173
		${}^1A_1'$	34.0	2.066	1.400	981
		${}^3A_1''$	33.0	2.287	1.307	1212
		${}^3A_1''$	33.0	2.287	1.307	1212
5	C_s	${}^1A_1'$	12.6 ^b	2.209	1.242	1291
		${}^3A_1''$	11.9	2.083	1.282	1220
6	C_s	${}^3A_1''$	8.3	2.661	1.202	1645
		${}^3A_1''$	8.3	2.661	1.202	1645

^a B3LYP/SSC(2g)/6-311++G(3dp,3df) single-point energies on B3LYP/SSC/6-311++G** optimized geometries.⁴ ^b This ground-state wave function has $M_s = 0$ (equal number of α - and β -spins), in which there is an α -spin on the U $5f_{\delta}$ orbital and a β -spin on the O₂ π^*_{yz} orbital. These orbitals overlap slightly, so the corresponding triplet state is slightly higher within the chosen scalar relativistic approximation. ^c An average of two U–O distances for the side-on (η^2) bound oxygen.

We propose that symmetric side-on binding of the superoxo species is favored because the 60° angle between adjacent lobes of the U $5f_{\phi}$ orbital gives an excellent overlap with adjacent lobes of the O₂ π^*_{xy} orbital, as shown in Figure 2. This contrasts with the case of d orbitals, where the 90° angle between adjacent lobes leads to a less favorable overlap with the O₂ π^*_{xy} orbital.⁷

The binding energy of dioxygen to U(V) systems is sensitive to the presence of electron-donating ligands that can increase the overall basicity of the uranium center, facilitating electron




Figure 2. Frontier MOs of UO_2^+ , O_2 , and their η^2 complex, showing a significant overlap between the uranyl $5f_{\phi}$ and O_2 π^*_{xy} orbitals.




Figure 3. B3LYP ligand binding energies for $[\text{UO}_2(\text{ACO})_n(\text{O}_2)]^+$ complexes (with and without zero-point energy (ZPE) corrections), where ACO = acetone and $n = 0–3$.

transfer to the O₂.³ This phenomenon is illustrated by trends in binding energies calculated for the $[\text{UO}_2(\text{ACO})_n(\text{O}_2)]^+$, where $n = 1–3$ and ACO = acetone. Coordination of a single ACO causes an increase in O₂ binding of ~ 5 kcal/mol compared to $[\text{UO}_2(\text{O}_2)]^+$, and addition of a second ACO results in a further strengthening of O₂ by another 4 kcal/mol (Figure 3). Addition of the third ACO does not appreciably change the O₂ binding energy. However, calculated ACO BE values decrease in a nearly linear fashion on going from $n = 1–3$ and become similar to the BE value for O₂ in the $[\text{UO}_2(\text{ACO})_3(\text{O}_2)]^+$ complex. Collision-induced dissociation studies³ showed that ACO is preferentially eliminated from $[\text{UO}_2(\text{ACO})_3(\text{O}_2)]^+$, while O₂ is lost from $[\text{UO}_2(\text{ACO})_2(\text{O}_2)]^+$. Thus, we expect the O₂ binding energies calculated with B3LYP to be slightly underestimated relative to the binding energies of acetone. (On the other hand, Table 2S and Figure 1S (Supporting Information) suggest that the LDA, PBE, and PW91 functionals yield O₂ binding energies that are too large. For example, in contrast to experiment,³ these methods predict O₂ to be more strongly bound than the second acetone ligand.)

The calculated U=O, U–O₂, and U–ACO bond lengths increase progressively with increasing n , consistent with normal behavior seen in metal ion coordination complexes.⁹ Interestingly, O–O bond lengths track the binding energy very closely, increasing as n goes from 0 to 2 and then decreasing modestly.

Figures 4 and 5 show the expected decrease of antisymmetric (ν_a) and symmetric (ν_s) O=U=O stretching frequencies with the number of ACO ligands coordinated to UO_2^+ and UO_2^{2+} . These results confirm that the U(VI) species have higher uranyl stretching frequencies than the corresponding U(V) species ($\Delta\nu_a = \sim 120$ cm⁻¹, $\Delta\nu_s = \sim 95$ cm⁻¹). The binding of dioxygen to U(V) dioxo cations results in a blue shift of uranyl vibrational frequencies. The frequency shift increases progressively from $\Delta\nu_a = 38$ cm⁻¹ and $\Delta\nu_s = 17$ cm⁻¹ to $\Delta\nu_a = 64$ cm⁻¹ and $\Delta\nu_s =$




Figure 4. Comparison of asymmetric O=U=O stretching frequencies for $[\text{UO}_2(\text{ACO})_n]^{2+}$, $[\text{UO}_2(\text{ACO})_n(\text{O}_2)]^+$, and $[\text{UO}_2(\text{ACO})_n]^+$ complexes.




Figure 5. Comparison of symmetric O=U=O stretching frequencies for $[\text{UO}_2(\text{ACO})_n]^{2+}$, $[\text{UO}_2(\text{ACO})_n(\text{O}_2)]^+$, and $[\text{UO}_2(\text{ACO})_n]^+$ complexes.

$= 42 \text{ cm}^{-1}$ as the number of acetone donor ligands increases from zero to three. This is consistent with the oxidation of U(V) to U(VI). Indeed, the O=U=O stretching modes for $\text{UO}_2^+\text{O}_2(\eta^2)$ ($\nu_a = 1039 \text{ cm}^{-1}$, $\nu_s = 947 \text{ cm}^{-1}$) are slightly higher than those for an unambiguous U(VI) compound, $(\text{UO}_2)^{2+}(\text{OH})^-$ ($\nu_a = 1017 \text{ cm}^{-1}$, $\nu_s = 937 \text{ cm}^{-1}$).

Our calculations also indicate that the presence of an additional 5f electron in the U(IV) complex (6), UOOH^+ (^3H state), further strengthens the interaction energy with dioxygen, leading to a bond energy of 36.3 kcal/mol (compared to 19.4 kcal/mol for the analogous U(V) complex 1). Here, the O_2 is again bound side-on with the O–O bond length of 1.32 Å and O–O stretching frequency of 1173 cm^{-1} . Again, we find a stable end-on bonding motif (complex 7), but it is 23.7 kcal/mol less stable. Interestingly, the reported^{5a} B3LYP binding energy for a neutral UO_2 (71.0 kcal/mol) is nearly twice as large as that for UOOH^+ (Table 1). Although no dioxygen–U(IV) complexes containing intact O_2 ligands have been observed, the products of irreversible oxidation to $[\text{UO}_2(\text{OH})]^+$ have been found in the gas phase.³

For the U(VI) complex $\text{UO}_2(\text{OH})^+$ 8, which lacks an available valence electron, we expect very weak binding to O_2 . Indeed,

TABLE 2: Natural Bond Order Analysis of Charge Transfer in 1–8^a

system	U	O ₂	O ₂ , α-spin	O ₂ , β-spin
1	0.191	−0.443	0.247	−0.690
2	−0.022	0.007	0.023	−0.015
3	0.156	−0.496	0.231	−0.727
4	0.133	−0.533	0.218	−0.751
5	0.097	−0.543	0.212	−0.755
6	0.234	−0.564	0.218	−0.782
7	0.145	−0.328	0.086	−0.414
8	−0.044	0.041	0.030	0.011

^a Obtained with B3LYP/SSC/6-311++G**.⁴

we find complex 8 to yield a $\text{U} \cdots \text{O}$ bond length of 2.66 Å and a bond strength of 8.3 kcal/mol. Since the standard entropy change upon complex formation is $-24.8 \text{ cal mol}^{-1} \text{ K}^{-1}$, this complex would not be expected to be observed at $T > 300 \text{ K}$. This is consistent with experiment.³

Natural bond order analysis⁸ of charge transfer in complexes 1–8 confirms (Table 2) that O_2 behaves as a weak electron donor in 2 and 8, whereas in 1, 3–6, and to some extent 7, it acts as a strong electron acceptor. In the latter case, there is almost full β-spin electron transfer (0.69–0.78e) from U to O_2 and notably large α-spin electron transfer in the opposite direction (0.22–0.25e), which is fully compatible with a superoxo character of the complexes.

In summary, our calculations suggest a novel two-electron three-atom bonding scheme for complexation of O_2 with U(V) compounds leading to the stabilization of superoxo complexes in the side-on (η^2) configuration. This binding motif, which is expected to be prevalent in many actinide compounds having valence $5f_\varphi$ electrons, is likely to play an important role in the oxidative processes involving these species.

Acknowledgment. Funding for this work was provided by the National Science Foundation (NIRT CTS Award # 0506951) and by the US Environmental Protection Agency (STAR Grant RD-83252501). This research was performed in part using the MSCF in EMSL, a national scientific user facility sponsored by the U.S. DOE, OBER and located at PNNL. W. A. de Jong acknowledges support from the BES Heavy Element Chemistry program of the U.S. Department of Energy, Office of Science.

Supporting Information Available: Table 1S provides additional information on geometric parameters and frequencies for U(IV), U(V), and U(VI) species and their complexes with O_2 . Table 2S and Figure 1S compare electronic binding energies and absolute energies for optimized structures at the B3LYP/SSC/6-311++G** level of theory. This material is available free of charge via the Internet at <http://pubs.acs.org>.

References and Notes

- (1) (a) Valentine, J. S. *Chem. Rev.* **1973**, *73*, 235. (b) Vaska, L. *Acc. Chem. Res.* **1976**, *9*, 175. (c) Reed, C. A.; Cheung, S. K. *Proc. Natl. Acad. Sci. USA* **1977**, *74*, 1780. (d) Momenteau, M.; Reed, C. A. *Chem. Rev.* **1994**, *94*, 659. (e) Girerd, J.-J.; Banse, F.; Simaan, A. J. Characterization and Properties of Non Heme Iron Peroxo Complexes. In *Metal-Oxo and Metal-Peroxo Species in Catalytic Oxidations*; Meunier, B., Ed.; Structure & Bonding; Springer: Berlin, 2000; Vol. 97, pp 145–177. (f) Borovik, A. C.; Zinn, P. G.; Zart, M. K. Dioxygen Binding and Activation: Reactive Intermediates. In *Activation of Small Molecules: Organometallic and Bioinorganic Perspectives*; Tolman, W. B., Ed; Wiley-VCH GmbH & Co. KGaA: New York, 2006; pp 187–234.
- (2) Bakac, A.; Espenson, J. H. *Inorg. Chem.* **1995**, *34*, 1730.

(3) Groenewold, G. S.; Cossel, K. C.; Gresham, G. L.; Gianotto, A. K.; Appelhans, A. D.; Olson, J. E.; Van Stipdonk, M. J.; Chien, W. *J. Am. Chem. Soc.* **2006**, *128*, 3075.

(4) Electronic structure calculations were carried out using the NWChem program.^{4a} Geometries were optimized at the B3LYP/SSC/6-311++G** level^{4b,c} followed by single point energy calculations using the Stuttgart-SC basis set^{4d} (SSC) augmented with the two g-functions of the Stuttgart-LC basis set on U and the 6-311++(3df,3pd) basis set on other elements. Control calculations for complex **1** show that this procedure leads to a small difference in binding energies (<0.5 kcal/mol) when compared to results obtained after full optimization at the B3LYP/SSC(3g,1h)/aug-cc-pVTZ level. A number of additional calculations were performed within the local density (LDA) and the generalized-gradient approximations (PBE^{4e} and PW91^{4f}). (a) Straatsma, T. P. et al. *NWChem, A Computational Chemistry Package for Parallel Computers, Version 5.0*; Pacific Northwest National Laboratory: Richland, WA 99354–0999, USA, 2006. (b) Becke, A. D. *Phys. Rev. A* **1988**, *38*, 3098. (c) Lee, C. T.; Yang, W. T.; Parr, R. G. *Phys. Rev. B* **1988**, *37*, 785. (d) Küsele, W.; Dolg, M.; Stoll, H.; Preuss, H. *J. Chem. Phys.* **1994**, *100*, 7535. (e) Perdew, J. P.; Burke, K.; Emzerhof, M. *Phys. Rev. Lett.* **1996**, *77*, 3865. (f) Perdew, J. P.; Wang, Y. *Phys. Rev. B* **1992**, *46*, 12947.

(5) The application of the B3LYP method for open-shell uranium complexes has proved a reasonable success.^{5a–c} For example, with the basis set used in this work, B3LYP yields the first (6.23 eV) and the second ionization potentials (15.22 eV) of UO₂ that are in reasonable agreement with experimental data:^{5f–i} 5.4–6.1 eV and 14.6–15.4 eV, respectively. (a) Zhou, M.; Andrews, L.; Ismail, N.; Marsden, C. *J. Phys. Chem. A* **2000**, *104*, 5495. (b) Majumdar, D.; Balasubramanian, K.; Nitsche, H. *Chem. Phys. Lett.* **2002**, *361*, 143. (c) Clavaguéra-Sarrio, C.; Vallet, V.; Maynau, D.; Marsden, C. *J. Chem. Phys.* **2004**, *121*, 5312. (d) Wang, X.; Andrews, L.; Li, J. *Inorg. Chem.* **2006**, *45*, 4157. (e) Shamov, G. A.; Schreckenbach, G.; Vo, T. N. *Chem.–Eur. J.* **2007**, *13*, 4932. (f) Cornehl, H. H.; Heinemann,

C.; Marcado, J.; Pires de Matos, A.; Schwarz, H. *Angew. Chem., Int. Ed. Engl.* **1996**, *35*, 891. (g) Allen, G. C.; Baerends, E. J.; Vernooijs, P.; Dyke, J. M.; Ellis, A. M.; Fehrer, M.; Morris, A. *J. Chem. Phys.* **1988**, *89*, 5363. (h) Han, J.; Goncharov, V.; Kaledin, L. A.; Komissarov, A. V.; Heaven, M. C. *J. Chem. Phys.* **2004**, *120*, 5155. (i) Gibson, J. K.; Haire, R. G.; Santos, M.; Marcado, J.; Pires de Matos, A. *J. Phys. Chem. A* **2005**, *109*, 2768.

(6) Pepper, M.; Bursten, B. E. *Chem. Rev.* **1991**, *91*, 719–741.

(7) Although the vast majority of known dioxygen–transition metal compounds are either in the end-on superoxo (η^1) or side-on peroxo (η^2) configuration, there have been some reports on η^2 1:1 complexes that can be better described as superoxides.^{7a–g} (a) Egan, J. W., Jr.; Haggerty, B. S.; Rheingold, A. L.; Theopold, K. H. *J. Am. Chem. Soc.* **1990**, *112*, 2445. (b) Fujisawa, K.; Tanaka, M.; Morooka, Y.; Kitajima, N. *J. Am. Chem. Soc.* **1994**, *116*, 12079. (c) Zhang, X.; Loppnow, G. R.; McDonald, R.; Takats, J. *J. Am. Chem. Soc.* **1995**, *117*, 7828. (d) Qin, K.; Incarvito, C. D.; Rheingold, A. L.; Theopold, K. H. *Angew. Chem., Int. Ed. Engl.* **2002**, *41*, 2333. (e) Spencer, D. J. E.; Aboeella, N. W.; Reynolds, A. M.; Holland, P. L.; Tolman, W. B. *J. Am. Chem. Soc.* **2002**, *124*, 2108. (f) Cramer, C. J.; Tolman, W. B.; Theopold, K. H.; Rheingold, A. L. *Proc. Natl. Acad. Sci. USA* **2003**, *100*, 3635. (g) Sarangi, R.; Aboeella, N.; Fujisawa, K.; Tolman, W. B.; Hedman, B.; Hodgson, K. O.; Solomon, E. I. *J. Am. Chem. Soc.* **2006**, *128*, 8286.

(8) (a) Glendenning, E. D.; Badenhop, J. K.; Reed, A. E.; Carpenter, J. E.; Bohmann, J. A.; Morales, C. M.; Weinhold, F. *NBO 5.50*; Theoretical Chemistry Institute, University of Wisconsin: Madison, WI, 2001; <http://www.chem.wisc.edu/~nbo5>. (b) Reed, A. E.; Curtiss, L. A.; Weinhold, F. *Chem. Rev.* **1988**, *88*, 899.

(9) Rodgers, M. T.; Armentrout, P. B. *Mass Spectrom. Rev.* **2000**, *19*, 215–247.

JP804202Q